

Anaerobic Organisms

Percentage susceptible

	Unasyn	Zosyn	Cefoxitin	Ertapenem	Imipenem	Meropenem	Clindamycin	Flagyl	Penicillin
Bacteroides spp.	78	94	70	82	98	95	33	100	–
Prevotella	–	100	–	–	100	98	69	99	100
Fusobacterium	100	96	–	–	95	100	77	95	–
Anaerobic Gram positive cocci*	–	99	–	–	99	100	97	100	100
Cutibacterium acnes**	–	100	–	–	94	–	53	0	–
Clostridiodes perfringens	100	100	–	–	100	100	83	100	90



Community Hospital
of the Monterey Peninsula
Montage Health

Reference: 2018 CLSI M-100, Performance Standards for Antimicrobial Susceptibility Testing, 28 th Edition.

* Anaerobic gram-positive cocci include Peptococcus, Peptostreptococcus, Finegoldia, Peptoniphilus and Anerococcus species

**Formerly Propionebacterium acnes

– No data available

Data were generated from unique isolates from patient specimens submitted to Tufts Medical center, Boston, Massachusetts; International Health Management Associates, Inc., Schaumburg, Illinois; R.M. Alden Research Laboratory Culver City, California; Creighton University School of medicine, Omaha, Nebraska; Mayo Clinic College of Medicine and Science, Rochester, Minnesota and the Centers for Disease Control and Prevention, Atlanta, Georgia. All testing was performed by the agar dilution method.