

# COVID-19 Vaccine Eligibility

Use this chart to help determine when to give COVID-19 vaccine based on age or health conditions.

Dose		Age:	5-11	12-15	16-17	18+
<b>Pfizer–Pediatric (5-11)</b>	<b>Primary 2 doses</b> (21 days apart)		✓			
<b>Pfizer–Adolescent/Adult (12+)</b>	<b>Primary 2 doses</b> (21 days apart)			✓	✓	✓
	<b>Additional (3rd) dose</b> for <a href="#">immunocompromised</a> , at least 28 days after 2nd Pfizer dose			✓	✓	✓
	<b>Booster dose*</b> of Pfizer (or different brand if age 18+), at least 6 months after 2nd dose of Pfizer				✓	✓
<b>Moderna (18+)</b>	<b>Primary 2 doses</b> (28 days apart)					✓
	<b>Additional (3rd) dose</b> for <a href="#">immunocompromised</a> , at least 28 days after 2nd Moderna dose					✓
	<b>Booster dose*</b> of Moderna ( <b>half-dose</b> ) or different brand, at least 6 months after 2nd dose of Moderna					✓
<b>Johnson &amp; Johnson (18+)</b>	<b>Primary 1 dose</b>					✓
	<b>Booster dose*</b> of J&J or different brand, at least 2 months after primary J&J dose					✓

\* Boosters are recommended for ages 16+ years, especially those at higher risk:

- Age 50+ years
- [Resident of long-term care facility](#)
- Underlying [medical condition](#) or increased risk of [social inequities](#)
- Risk of exposure from [work](#) or residence

View [Related Guidance and Requirements](#). (Eligibility is subject to change as guidance is updated.)